

Sortni poskus/ Variety trial

Rakičan

Leto/year 2017

Prejšnji posevek: ozimna pšenica

Gnojenje: 88 kg/ha N, 96 kg/ha P₂O₅, 120 kg/ha K₂O

Škropljenje: Successor 2,0 l/ha, Centium 0,25 l/ha, Revus 0,6 + 0,6 l/ha,

Pepelin 3,0 + 5,0 kg/ha, Pirimor 0,75 kg/ha, Mospilan 0,15 kg/ha

Gostota setve: medvrstna razdalja 140 cm, 1 kalivo seme/m²

Setev: 08.05.2017, spravilo: 13.09.2017

Zasnova poskusa: naključni blok v 4 ponovitvah, osnovna parcela 28 m²

Oljna buča - golica/ Oil pumpkin - hulles type

Cucurbita pepo

Previous crop: winter wheat

Fertilization: 88 kg/ha N, 96 kg/ha P₂O₅, 120 kg/ha K₂O

Spraying: Successor 2,0 l/ha, Centium 0,25 l/ha, Revus 0,6 + 0,6 l/ha,

Pepelin 3,0 + 5,0 kg/ha, Pirimor 0,75 kg/ha, Mospilan 0,15 kg/ha

Sowing density: interrow spacing 140 cm, 1 germinant seeds/m²

Sowing: 08.05.2017, harvesting: 13.09.2017

Trial layout: randomized block design in 4 repetitions, plot size 28 m²

SORTA VARIETY	Izvor Origin	Dobavitelj semena Seed supplier	Odpornost proti boleznim Resistance to disease	plodovi/ fruits				Absolutna masa zrnja Thousand seed weight	SM/CF		SB/CP		2017	
				nedozorele/immature	odpad/waste	uporabne/useful	skupaj/total		Vsebnost Contents	Pridelek Yield	Vsebnost Contents	Pridelek Yield	Zrnje s 9% vlage Grain with 9% moisture	Rodnostna skupina Yield category
			1-9 1=zelo dobra very good	%	%	%	število/ha number/ha	g	g/kg SS g/kg DM	kg/ha kg/ha	g/kg SS g/kg DM	kg/ha kg/ha	t/ha	
Gleisdorfer ölkurbis	A 2	SLO 2	3	5	7	88	17.054	255	448	732	363	593	1,80	I
Slovenska golica	SLO 3	SLO 3	3	9	8	83	14.196	213	451	505	360	403	1,23	II/3
Olinka	SRB 3	SLO 7	4	18	10	72	9.821	227	458	508	361	401	1,22	II/3
Povprečje/ Mean													1,41	
LSD (0,05)													0,25	