

Sortni poskus/Variety trial
Jablje

Prejšnji posevek: ozimna pšenica

Datum setve: 03.09.2010

Setvena količina: 20 kg/ha, medvrstna razdalja 15 cm

Zasnova poskusa: naključni blok v 4 ponovitvah, osnovna parcela 6 m²

Gnojenje v drugem letu rabe: 151 kg/ha N, 105 kg/ha P₂O₅, 105 kg/ha K₂O

Število košenj: 4 (10.05., 27.06., 14.08., 25.10.)

Navadna pasja trava/Cocksfoot
Dactylis glomerata

Previous crop: winter wheat

Date of sowing: 03.09.2010

Seed rate: 20 kg/ha, interrow spacing 15 cm

Trial layout: randomized block design in 4 repetitions, plot size 6 m²

Fertilization in second year: 151 kg/ha N, 105 kg/ha P₂O₅, 105 kg/ha K₂O

Number of cuttings: 4 (10.05., 27.06., 14.08., 25.10.)

SORTA VARIETY	Izvor Origin	Prezimitev Winter hardiness	Začetek latenja Heading date	Višina rastlin 1 košnje Plant height 1st cut	Intenzivnost latenja, 2 košnja Headind intensity, 2nd cut	Odpornost proti boleznim Resistance to disease	SB/CP		NEL/NEL		2012		2011-2012	
							Vsebnost Contents	Pridelek Yield	Vsebnost Contents	Pridelek Yield	Pridelek zelenja Herbage yield	Rodnostna skupina Yield category	Povprečni pridelek zelenja Average herbage yield	Povprečna relativna vrednost pridelka zelenja Average relative herbage yield
		1-9 1=zelo dobra very good	datum date	cm	1-9 1=brez without	1-9 1=zelo dobra very good	g/kg SS g/kg DM	kg/ha	MJ/kg SS MJ/kg DM	000 MJ/ha	t/ha (SS) t/ha (DM)		t/ha (SS) t/ha (DM)	%
DGŽ J01/03 30	SLO 1	1,0	6.5.	70	1,0	3,0	121	1545	5,67	72,6	12,81	II/1	11,20	104
Trerano	D 34	1,0	7.5.	65	1,0	4,0	116	1436	5,62	69,4	12,35	II/2	10,81	100
Kopa	SLO 1	1,0	6.5.	70	1,0	2,0	113	1359	5,69	68,5	12,05	II/3	10,44	96
Povprečje/Mean											12,40		10,82	100
LSD (0,05)											0,52			