

Krompir

Solanum tuberosum L.

V publikaciji predstavljamo rezultate posebnega preskušanja sort krompirja v letu 2018. Sortni poskusi so prikazani v dveh tematsko ločenih tabelah: pridelek in njegove karakteristike ter zdravstveno stanje z opazovanji razvojnih faz in opisi gomoljev. Dodane so še opombe, kjer so zapisane tudi v tabelah neopredeljene lastnosti. Glavnim tabelam so dodani podrobnejši rezultati o številu gomoljev. Opravljeni so bile tudi organoleptične ocene kakovosti kuhanega in ocvrtega krompirja. Vzorce je ocenjevala komisija v Poskusnem centru za krompir na IC Jablje Kmetijskega inštituta Slovenije v Mostah pri Komendi. Vzorci so bili pripravljeni po naslednjih postopkih:

- krompir je bil pred kuhanjem olupljen in kuhan na pari okoli 50 minut, čas kuhanja je bil odvisen od lastnosti sorte; opozoriti je še potrebno, da skupni vtis ni povprečna ocena vseh ocenjevanj temveč povsem samostojna ocena,
 - krompir smo cvrli pri 170°C v fritezi, čas cvrtja je bil odvisen od vsebnosti suhe snovi posamezne sorte.
- Prikazujemo še rezultate hitrosti polnjenja gomoljev - v šestih različnih terminih smo izkopali in stehtali po 5 zaporednih rastlin.

Poleg glavnega poskusa, kjer vrednotimo pridelek in njegove parametre, jedilno kakovost, vsebnost beljakovin in askorbinske kisline, skladiščenje in drugo, ugotavljamo tudi odpornost preskušanih sort na krompirjevo plesen, tolerantnost sort na herbicid Sencor, hitrost polnjenja gomoljev ter odpornost na virus Y^{NTN}. Ti podatki odločajo o primernosti sort za naše rastne razmere.

V letu 2018 je bilo v glavni sortni poskus vključenih 43 sort krompirja v Komendi in 26 sort v Rakičanu. Opravljeni so bile analize tal, pripravljeni tla so bila pogojena v skladu z analizami. Poskusni nasadi so bili posajeni po načrtu. Saditev je bila opravljena na vseh lokacijah v aprilu. Opravljeno je bilo škropljenje pred pleveli ter boleznimi in škodljivci. Opravljena je bila ocena bolezni poskusnih nasadov. Opravljena je bila ocena bolezni in vrednotenje fenofaz poskusnih nasadov. Opravili smo vzorčenje in izkop ter analize. Rastna sezona je bila za pridelovanje krompirja manj ugodna. Spomladi v času saditve je bilo precej mokro vreme, tako da je bila ta opravljena pozno. Tudi med rastno sezono je bilo zelo mokro vreme, tako da smo opazili veliko okužb s krompirjevo plesnijo. Pridelki na vseh lokacijah so bili zato povprečni. Opazili smo predvsem večje pokanje gomoljev. September je bil večji del sug, zato smo izkope opravili pravočasno.

Potato

The current publication contains the results of special testing of potato varieties performed in 2018. Variety trials are presented in two thematically separate tables: the yield and its characteristics and health condition with the monitoring of development phases and descriptions of tubers. Notes have been added in which properties undefined in tables are described. The main tables are accompanied by the more detailed results on the number of tubers. Organoleptic evaluation of the quality of cooked and fried potato was carried out. The samples were evaluated by the Commission at the Potato Centre at IC Jablje at Agricultural Institute of Slovenia in Moste near Komenda. The samples were prepared according to the following procedures:

- Prior to cooking the potato was peeled and vapour cooked for about 50 minutes, the time of cooking depended on the properties of varieties; it has to be noted that the overall impression is not an average value of all evaluations but an entirely independent one,
- The potato was fried at 170°C in a frying pan, the time of frying depended on the content of dry matter of individual varieties.

The results of the tuber bulking rate are presented – on six different dates 5 consecutive plants were excavated and weighed.

Beside the main trial in which the yield and its parameters, edible quality, protein and ascorbic acid content, storage and other properties are evaluated, the resistance of tested varieties to potato late blight, the tolerance of varieties to the herbicide Sencor, tuber bulking rate and the resistance to virus Y^{NTN} are determined. These data decide on the suitability of varieties for the Slovene growing conditions.

43 potato varieties were grown in trials in Komenda and 26 varieties in Rakičan respectively in 2018. Soil analyzes were carried out, the prepared soil was fertilized according to the analyzes. Experimental crops were planted according to plan. Planting was carried out at all locations in April. Spraying for weeds and diseases and pests was done on time. An assessment of the disease of experimental crops was carried out. An evaluation

of the disease and the evaluation of the pheno- phases of the experimental crops were carried out. The samples were excavated and analyzed in October. The growing season was less favorable for the cultivation of potatoes. In the spring, during the planting, it was wet, so it was done late. The summer was also wet, so there were a lot of problems with late blight. Therefore there were average yields at all locations. The tuber cracks were observed on tuber surface at several varieties. September was very dry, so we harvested trials on time.